



TITLE:

# Mediterranean mimicker

AUTHOR(S):

Arasawa, Soichi; Nakase, Hiroshi; Ozaki, Yoshinao;  
Uza, Norimitsu; Matsuura, Minoru; Chiba, Tsutomu

---

CITATION:

Arasawa, Soichi ...[et al]. Mediterranean mimicker. The Lancet 2012,  
380(9858): 2052-2052

ISSUE DATE:

2012-12

URL:

<http://hdl.handle.net/2433/167317>

RIGHT:

© 2012 Elsevier Ltd.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Arasawa *et al*

## **Familial Mediterranean Fever with Colonic Involvement Mimicking Inflammatory**

### **Bowel Disease**

Soichi Arasawa, MD, Hiroshi Nakase, MD, Yoshinao Ozaki, MD, Norimitsu Uza, MD,

Minoru Matsuura, MD, Tsutomu Chiba, MD

Department of Gastroenterology and Hepatology, Graduate School of Medicine, Kyoto  
University.

Corresponding to: Hiroshi Nakase, MD,

Department of Gastroenterology and Hepatology, Graduate School of

Medicine, Kyoto University

54 Shogoin-Kawaharacho, Sakyo-ku, Kyoto-city,

Kyoto, 606-8507, Japan

Tel: +81-75-751-4319      Fax: +81-75-751-4303

E-mail: [hiropy\\_n@kuhp.kyoto-u.ac.jp](mailto:hiropy_n@kuhp.kyoto-u.ac.jp)

Word Count: 599 words

A 42-year-old woman experienced recurrent febrile attacks (over 38°C) and diffuse abdominal pain that spontaneously subsided within two to three days, beginning in April 2009.

Laboratory data showed leukocytosis and elevated serum C-reactive protein. Colonoscopy revealed discontinuous loss of fine vascular markings, erythema, and friable mucosa in the right hemicolon. Biopsy specimens revealed no characteristic features related to Crohn's disease (CD) or ulcerative colitis (UC). CT scan showed wall thickness of colon. Capsule endoscopy revealed no abnormality in small intestine. ASCA and ANCA were negative.

Therefore, she was diagnosed with indeterminate colitis. 4g of mesalazine was firstly administered, and 1.5g was kept for maintenance. But her symptoms continued.

In July 2009, 40mg of prednisolone was orally administered to control her repeated episodes.

Her symptoms subsided, and the prednisolone dosage was tapered in August 2009. Her symptoms recurred, however, when the prednisolone dosage reached the 6-mg level.

Mercaptopurine could not be used because of her intolerability.

In April 2010, colonoscopy revealed circumferentially erythematous mucosa with erosions in the cecum, longitudinal erosions with pseudopolypoid-like lesions in the right hemicolon, and no involvement of the rectum. These endoscopic features resemble CD (Figure 1-A and

1-C), but biopsy specimens revealed no granuloma. Intestinal tuberculosis and other infectious diseases were ruled out.

She continued to have febrile attacks every week in May 2010. Moreover, her attacks were accompanied by severe abdominal pain as well as bilateral arthralgia in the wrists, elbows, and shoulders.

Because her symptoms met the Tel Hashomer criteria, we suspected familial Mediterranean fever (FMF). After obtaining written informed consent, we performed a genome sequencing examination for the Mediterranean fever gene (*MEFV*) mutation, which revealed a heterozygous *MEFV* mutation, a 910G>A point mutation. This mutation causes a Gly304Arg missense mutation on pyrin (Figure 2).

After diagnosing her with FMF in late May 2010, 1mg of colchicine was administered. Her symptoms promptly subsided, and we could stop prednisolone and mesalazine. In February 2011, colonoscopy revealed remarkable improvement of the mucosal inflammation in the colon (Figure 1-B and 1-D). In the two years since starting colchicine treatment, she has had no febrile attacks or serositic symptoms.

FMF is a periodic inflammatory disease characterized by episodic febrile attacks and serositis.

It is a monogenic Mendelian disease caused by *MEFV* on chromosome 16p13.3, which encodes pyrin.<sup>1</sup> Pyrin is a 781-amino-acid protein mainly expressed in granulocytes and monocytes,<sup>2</sup> and regulates the inflammatory response by blocking intracellular signal pathways via NF- $\kappa$ B or caspase 1. The absence of pyrin function due to mutated *MEFV* leads to the oversecretion of inflammatory cytokines, resulting in FMF.<sup>3</sup> FMF is rare in non-Mediterranean populations, which might mislead physicians to diagnose inflammatory bowel disease (IBD) in patients who present with the intestinal symptoms of FMF. Recent studies suggest an association between *MEFV* and IBD,<sup>4</sup> although *MEFV* mutations make no significant contribution to IBD susceptibility.<sup>5</sup> This patient with FMF showed colonic lesions mimicking CD. Despite her refractoriness to conventional IBD treatment, colchicine dramatically relieved her symptoms with the improvement of colonic lesions, and she has had no relapse in the last 2 years. This clinical course and genetic analysis led to a diagnosis of colonic lesions related to FMF. Local circulatory disturbance at colonic wall by sustained inflammation related to FMF might result in mucosal damage. Our case suggests that some patients with FMF might have IBD-like mucosal lesions and physicians should consider genetic analysis and colchicine treatment for patients with indeterminate colitis that

do not respond to conventional IBD treatment.

### **Contributors**

SA, HN, YO, and NU looked after the patient. SA, HN, MM, and TC wrote the report.

Written consent to publish was obtained.

## References

1. Berdeli A, Mir S, Nalbantoglu S, et al. Comprehensive Analysis of a Large-Scale Screen for MEFV Gene Mutations: Do They Truly Provide a “Heterozygote Advantage” in Turkey? *Genet Test Mol Biomarkers*. 2011; 15(7-8): 475-82.
2. Centola M, Wood G, Frucht D, et al. The gene for familial Mediterranean fever, MEFV, is expressed in early leukocyte development and is regulated in response to inflammatory mediators. *Blood*. 2000; 95 (10): 3223-31.
3. Ting JP, Kastner DL, Hoffman HM. CATERPILLERS, pyrin and hereditary immunological disorders. *Nat Rev Immunol*. 2006; 6(3): 183-95.
4. Yıldırım B, Tuncer C, Kan D, et al. MEFV gene mutations and its impact on the clinical course in ulcerative colitis patients. *Rheumatol Int*. 2011; 31(7): 859-64.
5. Villani A, Lemire M, Louis E, et al. Genetic variation in the familial Mediterranean fever gene (MEFV) and risk for Crohn's disease and ulcerative colitis. *PLoS One*. 2009; 4(9): e7154.

## Figure Legends

**Figure 1.** Colonoscopic views at the cecum and transverse colon before (A, C) and after (B, D) colchicine treatment.

A: Circumferentially erythematous mucosa with erosions at cecum. B: Remarkable improvement of colonic inflammation with patchy erosions at cecum. C. Pseudopolypoid like lesions at transverse colon. D. Scar formation alone at transverse colon.

**Figure 2.** Genetic analysis for the Mediterranean fever gene mutation. A heterozygous mutation in *MEFV* gene, a 910G>A point mutation, was revealed. This genetic mutation causes a Gly304Arg missense mutation on pyrin. The sequencing number was counted from the head of the primer.